

Effect of a mobilization and active exercise program on the range of motion of bedridden patients with disuse syndrome

Teresa Fernandes, Eugénia Mendes, Leonel Preto, André Novo
School of Health, Polytechnic Institute of Bragança – Portugal



Keywords: Disuse Syndrome; Goniometer; Articular mobilization; Active exercise || maria.mendes@ipb.pt

INTRODUCTION

Disuse syndrome is a disorder that is most often associated with acute or chronic disease complications. Despite disuse syndrome may affect all organs and systems, the impact of range of motion limitations caused by immobility on functional capacity to perform activities of daily living is often very severe. Mobilization and active exercise have beneficial effects that counteract the impact of immobility on the body.

OBJECTIVE

This study aims to assess the effect of a mobilization and active exercise program on the range of motion of bedridden patients with disuse syndrome.

METHOD

A quasi-experimental study was developed. The sample consisted of 26 persons that have been bedridden for more than six months at home. A mobilization and active exercise program was design, fitting patient individual needs and implemented 2 times/week for 2 months. Caregivers where trained to transfer the patient from bed to chair and to repeat active exercise every day. Data collection was performed before and after intervention, using the Barthel Index and a goniometer for range of motion evaluation.



RESULTS

26 participants, aged 77.19 ± 11.67 and bedridden for 18 months (18.73 ± 15.25) were enrolled, but only 24 completed the intervention program. There was a minimal difference in the sex distribution with 7.6% more women than men. Results showed a statistically significant increase on range of motion of the shoulder, elbow, wrist, hip and knee. There was statistical significance in plantar flexion but not on the dorsiflexion. Barthel Index score has increase significantly ($28,65 \pm 21,28$ vs $31,46 \pm 23,28$; $p=0,035$) after the mobilization and active exercise program.

	Left Shoulder				Rigth Shoulder			
	Abduction	Abduction	Flexion	Flexion	Abduction	Abduction	Flexion	Flexion
	T0	T1	T0	T1	T0	T1	T0	T1
N	24	24	24	24	24	24	24	24
Mean	50,77	56,67	53,46	59,58	49,42	55	55,77	61,67
Standard Deviation	47,99	51,38	51,33	53,78	46,67	49,95	49,71	51,59
Z	-3,581		-3,758		-3,463		-3,463	
P	0		0		0,001		0,001	

	Left Elbow		Rigth Elbow	
	Flexion	Flexion	Flexion	Flexion
	T0	T1	T0	T1
N	24	24	24	24
Mean	65,96	71,04	65,96	72,29
Standard Deviation	39,01	40,72	38,41	39,12
Z	-3,945		-3,906	
P	0		0	

	Left Wrist				Rigth Wrist			
	Palmar Flexion	Palmar Flexion	Dorsal Flexion	Dorsal Flexion	Palmar Flexion	Palmar Flexion	Dorsal Flexion	Dorsal Flexion
	T0	T1	T0	T1	T0	T1	T0	T1
N	24	24	24	24	24	24	24	24
Mean	31,92	35,21	27,12	30,21	31,35	35,63	26,92	29,58
Standard Deviation	18,55	20,51	16,86	18,27	17,41	18,84	16,25	17,5
Z	-3,3		-3,207		-3,945		-2,887	
p	0,001		0,001		0		0,004	

CONCLUSION

A mobilization and active exercise program implemented regularly may contribute to improve range of motion of bedridden with disuse syndrome. Thus, based on these results, it is believed that the intervention of rehabilitation nurse was an added value, contributing to an increase in range of motion and independence of patients.

	Left Hip				Rigth Hip			
	Abduction	Abduction	Flexion	Flexion	Abduction	Abduction	Flexion	Flexion
	T0	T1	T0	T1	T0	T1	T0	T1
N	24	24	24	24	24	24	24	24
Mean	17,88	20,63	29,81	35	17,31	20,83	29,04	33,54
Standard Deviation	10,11	11,63	18,57	19,22	9,51	10,49	13,93	14,25
Z	-3,3		-3,879		-3,69		-4,123	
p	0,001		0		0		0	

	Left Knee		Rigth Knee	
	Flexion	Flexion	Flexion	Flexion
	T0	T1	T0	T1
N	24	24	24	24
Mean	60,58	67,71	61,92	70
Standard Deviation	32,44	32,06	30,66	31,24
Z	-3,841		-4,062	
P	0		0	

REFERENCES

Fan, E., Zanni, J. M., Dennison, C. R., Lepre, S. J., & Needham, D. M. (2009). Critical Illness Neuromyopathy and Muscle Weakness in Patients in the Intensive Care Unit. *AACN Advanced Critical Care*, 243-253.

Heer, M., Baecker, N., Mika, C., Boese, A., & Gerzer, R. (2005). Immobilization induces a very rapid increase in osteoclast activity. *Acta Astronautica* 57, 31-36

Lambert, C. P. & Evans, W. J. (2002). Effects of Aging and Resistance Exercise on Determinants of Muscle Strength. *Journal of the American Aging Association* 25, 73-78.

Morris, P. E., Goad, A., Thompson, C., Taylor, K., Harry, B., Passmore, L., Ross, A., ... Haponik, E. (2008). Early intensive care unit mobility therapy in the treatment of acute respiratory failure. *Critical Care Medicine*, 36 (8), 2238-2243.

Netz, Y., Wu, M., Becker, B. J. & Tenenbaum, G. (2005). Physical Activity and Psychological Well-Being in Advanced Age: A Meta-Analysis of Intervention Studies. *Psychology and Aging* 20 (2), 272-284.

Perme, C., Southard, R., Joyce, D., Noon, G. & Loebe, M. (2006). Early mobilization of LVAD recipients who require prolonged mechanical ventilation. *Texas Heart Institute Journal*, 33(2), 130-133.

	Left Foot				Rigth Foot			
	Plantar Flexion	Plantar Flexion	Dorsal Flexion	Dorsal Flexion	Plantar Flexion	Plantar Flexion	Dorsal Flexion	Dorsal Flexion
	T0	T1	T0	T1	T0	T1	T0	T1
N	24	24	24	24	24	24	24	24
Mean	17,5	19,79	7,69	8,75	18,08	21,04	7,69	8,33
Standard Deviation	11,42	13,86	4,29	5,56	11,05	13,1	3,53	4,34
Z	-2,496		-2,236		-3		-1,732	
p	0,013		0,025		0,003		0,083	

	Barhtel Index T0	Barhtel Index T1
N	24	24
Mean	28,65	31,46
Standard Deviation	21,28	23,28
Z	-2,111	
p	0,035	